

Anti-JNK Rabbit pAb



WL01295

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-JNK Rabbit pAb		
Source	Rabbit		
Species reactivity	Human, Mouse, Rat		
Tested applications	WB	1:500-1:1000	
	IHC	1:150-1:300	
Cellular localization	Secreted and Cell membrane		
Pack size	50/100/200/500/1000µl		
Storage	Store at -20°C. Avoid freeze/thaw cycles.		
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide		

General Information

Background

c-Jun N-terminal kinases (JNKs) phosphorylate and augment transcriptional activity of c-Jun. JNKs originate from three genes that yield 10 isoforms through alternative mRNA splicing, including JNK1a1, JNK1b1, JNK2a1, JNK2b1, and JNK3a1, which represent the p46 isoforms, and JNK1a2, JNK1b2, JNK2a2, JNK2b2, and JNK3b2, which represent the p54 isoforms. JNKs coordinate cell responses to stress and influence regulation of cell growth and transformation. The human JNK1 (PRKM8, SAPK1, MAPK8) gene maps to chromosome 10q11.22 and shares 83% amino acid identity with JNK2. JNK1 is necessary for normal activation and differentiation of CD4 helper T (TH) cells into TH1 and TH2 effector cells. Capsaicin activates JNK1 and p38 in ras-transformed human breast epithelial cells. Nitrogen oxides (NOx) upregulate JNK1 in addition to c-Fos, c-Jun, and other signaling kinases, including MEKK1 and p38. Polyclonal antibody is produced by immunizing animals with a synthetic peptide of JNK.

Immunogen

Purification

Polyclonal antibody was purified by immunogen affinity chromatography.


Anti-JNK Rabbit pAb



WL01295


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Western blot-Anti-JNK pAb

Lane 1: Human HeLa cell lysate
 Lane 2: Human MCF-7 cell lysate
 Lane 3: Human HEK-293 cell lysate
 Lane 4: Human MGC-803 cell lysate
 All lanes: Anti-JNK at 1:1000 dilution
 Lysates/proteins at 20-50 µg per lane.
 Predicted band size: 46,54 kDa
 Observed band size: 46,54 kDa



Immunohistochemistry-Anti-JNK pAb

Immunohistochemical analysis of paraffin-embedded human pancreatic cancer using anti-JNK Rabbit Antibody at 1:150 dilution.
 Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunohistochemistry-Anti-JNK pAb

Immunohistochemical analysis of paraffin-embedded rat colon using anti-JNK Rabbit Antibody at 1:300 dilution.
 Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0