Wanleibio

Anti-Frizzled 8 Rabbit pAb

WL04715

For Research Use Only.Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-Frizzled 8 Rabbit pAb	
Source	Rabbit	
Species reactivity	Human	
Tested applications	WB	1:500-1:1000
Packsize	50/100/200/500/1000µl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 $\mu g/ml$	
	BSA, 50% glycerol and less than 0.02% sodium azide	


Anti-Frizzled 8 Rabbit pAb

Wanleibio

WL04715

For Research Use Only.Not For Use In Diagnostic Procedures

Product Images



Western blot-Anti-Frizzled 8 pAb

Lane 1: Human Hela cell lysate Lane 2: Human MCF-7 cell lysate Lane 3: Human HEK-293 cell lysate Lane 4: Human A549 cell lysate All lanes: Anti-Frizzled 8 at 1:1000 dilution Lysates/proteins at 20-50 µg per lane. Predicted band size: 73 kDa Observed band size: 73 kDa

General Information

BackgroundIntronless gene is a member of the frizzled gene family. Members of this
family encode seven-transmembrane domain proteins that are receptors
for the Wingless type MMTV integration site family of signaling proteins.
Most frizzled receptors are coupled to the beta-catenin canonical signaling
pathway. This gene is highly expressed in two human cancer cell lines,
indicating that it may play a role in several types of cancer.

Immunogen	Polyclonal antibody is produced by immunizing animals with a synthetic peptide of Frizzled 8.
Purification	Polyclonal antibody was purified by immunogen affinity chromatography.