

Anti-NK-p44 Rabbit pAb



WL05307

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-NK-p44 Rabbit pAb	
Source	Rabbit	
Species reactivity	Human	
Tested applications	WB	1:1000-1:2000
Pack size	50/100/200/500/1000μl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide	

General Information

Background	Cd336, also named as NCR2, LY95 and Nkp44, belongs to the natural cytotoxicity receptor (NCR) family. It is cytotoxicity-activating receptor that may contribute to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis. It is expressed on activated human NK cells. CD336 displays a single extracellular Ig-like V domain and a transmembrane portion containing the charged residue (Lysine), likely involved in the association with KARAP/DAP12 molecules.
Immunogen	Polyclonal antibody is produced by immunizing animals with a synthetic peptide of NK-p44.
Purification	Polyclonal antibody was purified by Protein A affinity chromatography.


Anti-NK-p44 Rabbit pAb



WL05307

For Research Use Only. Not For Use In Diagnostic Procedures

Product Images



Western blot-Anti-NK-p44 pAb

Lane 1: Human MCF-7 cell lysate

Lane 2: Human HEK-293 cell lysate

All lanes: Anti-NK-p44 at 1:1000 dilution

Lysates/proteins at 20-50 μg per lane.

Predicted band size: 31 kDa

Observed band size: 31 kDa