

Anti-IL-17RA Rabbit pAb



WL04433

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-IL-17RA Rabbit pAb
Source	Rabbit
Species reactivity	Human, Mouse, Rat
Tested applications	WB 1:2000
Cellular localization	Secreted and Cell membrane
Pack size	50/100/200/500/1000µl
Storage	Store at -20°C. Avoid freeze/thaw cycles.
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide

General Information

Background Interleukin 17A (IL17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure.

Immunogen Polyclonal antibody is produced by immunizing animals with a synthetic peptide of IL-17RA.

Purification Polyclonal antibody was purified by Protein A affinity chromatography.


Anti-IL-17RA Rabbit pAb



WL04433


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images



Western blot-Anti-IL-17RA pAb

Lane 1: Human HeLa cell lysate
 Lane 2: Human HEK-293 cell lysate
 Lane 3: Human A549 cell lysate
 All lanes: Anti-IL-17RA at 1:2000 dilution
 Lysates/proteins at 20-50 µg per lane.
 Predicted band size: 96 kDa
 Observed band size: 96-120 kDa



Western blot-Anti-IL-17RA pAb

Lane 1: Mouse heart tissue lysate
 Lane 2: Mouse liver tissue lysate
 All lanes: Anti-IL-17RA at 1:2000 dilution
 Lysates/proteins at 20-50 µg per lane.
 Predicted band size: 96 kDa
 Observed band size: 96-120 kDa