

Anti-APE1/Ref-1 Rabbit pAb



WL02884

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-APE1/Ref-1 Rabbit pAb	
Source	Rabbit	
Species reactivity	Human, Mouse, Rat	
Tested applications	WB	1:500-1:1000
	IHC	1:100-1:400
Cellular localization	Cytoplasm;Endoplasmic reticulum;Mitochondrion;Nucleus	
Pack size	50/100/200/500/1000µl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide	

General Information

Background	<p>Apurinic/aprimidinic (AP) sites occur frequently in DNA molecules by spontaneous hydrolysis, by DNA damaging agents or by DNA glycosylases that remove specific abnormal bases. Ape1 (Apurinic/Apyrimidic eEndonuclease 1), also known as Ref1 (Redox effector factor 1), is a multifunctional protein with several biological activities. These include roles in DNA repair and in the cellular response to oxidative stress. Ape1 initiates the repair of abasic sites and is essential for the base excision repair (BER) pathway. Repair activities of Ape1 are stimulated by interaction with XRCC1, another essential protein in BER. Ape1 functions as a redox factor that maintains transcription factors in an active, reduced state but can also function in a redox-independent manner as a transcriptional cofactor to control different cellular fates such as apoptosis, proliferation and differentiation.</p>
Immunogen	<p>Polyclonal antibody is produced by immunizing animals with a synthetic peptide of APE1/Ref-1.</p>

Purification Polyclonal antibody was purified by immunogen affinity chromatography.


Anti-APE1/Ref-1 Rabbit pAb



WL02884


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Western blot-Anti-APE1/Ref-1 pAb

Lane 1: Human HepG2 cell lysate
 Lane 2: Human Hela cell lysate
 Lane 3: Human BGC-823 cell lysate
 Lane 4: Human MGC-803 cell lysate
 All lanes: Anti-APE1/Ref-1 at 1:1000 dilution
 Lysates/proteins at 20-50 µg per lane.
 Predicted band size: 35 kDa
 Observed band size: 37 kDa



Western blot-Anti-APE1/Ref-1 pAb

Lane 1: Mouse kidney tissue lysate
 Lane 2: Mouse heart tissue lysate
 Lane 3: Rat brain tissue tissue lysate
 All lanes: Anti-APE1/Ref-1 at 1:1000 dilution
 Lysates/proteins at 20-50 µg per lane.
 Predicted band size: 35 kDa
 Observed band size: 37 kDa



Immunohistochemistry-Anti-APE1/Ref-1 pAb

Immunohistochemical analysis of paraffin-embedded mouse lung using anti-APE1/Ref-1 Rabbit Antibody at 1:150 dilution.
 Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0