

Anti-Bax Rabbit pAb



WL01637

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-Bax Rabbit pAb		
Source	Rabbit		
Species reactivity	Human, Rat, Mouse, Rabbit		
Tested applications	WB	1:1000-1:2000	
	IHC	1:300	
	IF	1:300-1:500	
Cellular localization	Cytoplasm and Mitochondrion membrane		
Pack size	50/100/200/500/1000μl		
Storage	Store at -20°C. Avoid freeze/thaw cycles.		
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 μg/ml		
	BSA, 50% glycerol and less than 0.02% sodium azide		

General Information

Background	The Bax encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apoptotic activator. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis.
Immunogen	Polyclonal antibody is produced by immunizing animals with a synthetic peptide of Bax.
Purification	Polyclonal antibody was purified by protein A affinity chromatography.


Anti-Bax Rabbit pAb



WL01637


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Western blot-Anti-Bax pAb

Lane 1: Human HepG2 cell lysate
 Lane 2: Human Hela cell lysate
 Lane 3: Human BGC-823 cell lysate
 Lane 4: Human MGC-803 cell lysate
 All lanes: Anti-Bax at 1:1000 dilution
 Lysates/proteins at 20-50 μg per lane.
 Predicted band size: 23 kDa
 Observed band size: 23 kDa



Western blot-Anti-Bax pAb

Lane 1: Mouse kidney tissue lysate
 Lane 2: Mouse heart tissue lysate
 Lane 3: Rat liver tissue lysate
 Lane 4: Rat brain lysate
 All lanes: Anti-Bax at 1:1000 dilution
 Lysates/proteins at 20-50 μg per lane.
 Predicted band size: 23 kDa
 Observed band size: 23 kDa



Western blot-Anti-Bax pAb

Lane 1: Rabbit heart tissue lysate
 Lane 2: Rabbit kidney tissue lysate
 Lane 3: Rabbit liver tissue lysate
 Lane 4: Rabbit stomach tissue lysate
 All lanes: Anti-Bax at 1:1000 dilution
 Lysates/proteins at 20-50 μg per lane.
 Predicted band size: 23 kDa
 Observed band size: 23 kDa


Anti-Bax Rabbit pAb



WL01637


For Research Use Only. Not For Use In Diagnostic Procedures

Product Information




Immunohistochemistry-Anti-Bax pAb

Immunohistochemical analysis of paraffin-embedded human brain using anti-Bax Rabbit Antibody at 1:250 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunohistochemistry-Anti-Bax pAb

Immunohistochemical analysis of paraffin-embedded human colon cancer using anti-Bax Rabbit Antibody at 1:500 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunohistochemistry-Anti-Bax pAb

Immunohistochemical analysis of paraffin-embedded mouse liver using anti-Bax Rabbit Antibody at 1:250 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0


Anti-Bax Rabbit pAb



WL01637


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Immunohistochemistry-Anti-Bax pAb

Immunohistochemical analysis of paraffin-embedded rat colon using anti-Bax Rabbit Antibody at 1:300 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunohistochemistry-Anti-Bax pAb

Immunohistochemical analysis of paraffin-embedded rabbit colon using anti-Bax Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0




Immunohistochemistry-Anti-Bax pAb

Immunohistochemical analysis of paraffin-embedded rabbit submaxillary glands using anti-Bax Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0


Anti-Bax Rabbit pAb

For Research Use Only. Not For Use In Diagnostic Procedures


Product Information



Immunohistochemistry-Anti-Bax pAb
Immunohistochemical analysis of paraffin-embedded rabbit adrenal gland using anti-Bax Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunohistochemistry-Anti-Bax pAb
Immunohistochemical analysis of paraffin-embedded rabbit uterus using anti-Bax Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunofluorescence-Anti-Bax pAb
Immunofluorescence analysis of paraffin-embedded Human colon cancer using anti-Bax Rabbit Antibody at 1:500 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0




WL01637


Anti-Bax Rabbit pAb

For Research Use Only. Not For Use In Diagnostic Procedures

Product Images



Immunofluorescence-Anti-Bax pAb
Immunofluorescence analysis of paraffin-embedded mouse lung using anti-Bax Rabbit Antibody at 1:300 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunofluorescence-Anti-Bax pAb
Immunofluorescence analysis of paraffin-embedded rat lung using anti-Bax Rabbit Antibody at 1:500 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



WL01637