Product Datasheet

Wanleibio

Anti-IL6 Rabbit pAb



WL02841

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name Anti-IL6 Rabbit pAb

Source Rabbit

Human, Mouse, Rat **Species reactivity**

Tested applications WB 1:1000-1:2000

> IHC 1:100-1:400

ΙF 1:300

Cellular localization Secreted

50/100/200/500/1000µl Pack size

Store at -20°C. Avoid freeze/thaw cycles. **Storage**

Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml Storage buffer

BSA, 50% glycerol and less than 0.02% sodium azide

General Information

Background

IL6(Interleukin-6) is a multifunctional protein, that plays critical roles in host defense, immune response and hematopoiesis. IL6 is constitutively expressed by epidermal Langerhans cells and its expression is induced in stimulated keratinocytes. IL6 induces proliferation or differentiation in many cell types including B cells, thymocytes and T cells. IL6, in concert with TGF-β, is important for developing Th17 responses. IL6 binds to IL6Rα and through this association induces gp130 homodimerization. Research studies have shown that IL6, through increasing expression of proangiogenic VEGF, may also contribute to metastatic breast cancer.

Immunogen Polyclonal antibody is produced by immunizing animals with a synthetic

peptide of IL6.

Purification Polyclonal antibody was purified by Protein A affinity chromatography.

Product Datasheet

Wanleib




WL02841

Anti-IL6 Rabbit pAb


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Western blot-Anti-IL6 pAb

Lane 1: Human HepG2 cell lysate Lane 2: Human Hela cell lysate Lane 3: Human BGC-823 cell lysate Lane 4: Human MGC-803 cell lysate All lanes: Anti-IL6 at 1:1000 dilution Lysates/proteins at 20-50 µg per lane. Predicted band size: 24 kDa Observed band size: 21 kDa



Western blot-Anti-IL6 pAb

Lane 1: Mouse kidney tissue lysate Lane 2: Mouse brain tissue lysate Lane 3: Rat spleen tissue lysate Lane 4: Rat stomach tissue lysate All lanes: Anti-IL6 at 1:1000 dilution Lysates/proteins at 20-50 µg per lane. Predicted band size: 24 kDa Observed band size: 21 kDa



Immunohistochemistry-Anti-IL6 pAb

Immunohistochemical analysis of paraffin-embedded rat lung using anti-IL6 Rabbit Antibody at 1:150 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0


Wanleibio Co., Ltd. 400-602-0407 Wanleibio Co., Ltd. 400-602-0407 www.wanleibio.com sales@wanleibio.com www.wanleibio.com sales@wanleibio.com

Anti-IL6 Rabbit pAb



For Research Use Only. Not For Use In Diagnostic Procedures


Product Images



Immunohistochemistry-Anti-IL6 pAb

Immunohistochemical analysis of paraffin-embedded mouse colon using anti-IL6 Rabbit Antibody at 1:300 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunofluorescence-Anti-IL6 pAb

Immunofluorescence analysis of paraffin-embedded mouse lung using anti-IL6 Rabbit Antibody at 1:300 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

Wanleibio Co.,Ltd. 400-602-0407 sales@wanleibio.com www.wanleibio.com

