

Anti-SMAD3 Rabbit pAb



WL02288

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-SMAD3 Rabbit pAb	
Source	Rabbit	
Species reactivity	Human, Mouse, Rat	
Tested applications	WB	1:1000-1:2000
	IHC	1:200
	IF	1:300
Cellular localization	Cytoplasm and Nucleus	
Pack size	50/100/200/500/1000µl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide	

General Information

Background	Members of the Smad family of signal transduction molecules are components of a critical intracellular pathway that transmit TGF-β signals from the cell surface into the nucleus. Smad1 and Smad5 are effectors of BMP-2 and BMP-4 function, while Smad2 and Smad3 are involved in TGF-β and Activin-mediated growth modulation. Smad4 has been shown to mediate all of the above activities through interaction with various Smad family members. The phosphorylated receptor-regulated Smad associates from the receptor and forms a heteromeric complex with the co-Smad (Smad4), allowing translocation of the complex to the nucleus. Once in the nucleus, Smads can target a variety of DNA binding proteins to regulate transcriptional responses.
Immunogen	Polyclonal antibody is produced by immunizing animals with a synthetic peptide of SMAD3.
Purification	Polyclonal antibody was purified by Protein A affinity chromatography.


Anti-SMAD3 Rabbit pAb



WL02288


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Western blot-Anti-SMAD3 pAb

Lane 1: Human BGC-823 cell lysate
 Lane 2: Human MGC-803 cell lysate
 Lane 3: Human SGC-7901 cell lysate
 Lane 4: Human MCF-7 cell lysate
 All lanes: Anti-SMAD3 at 1:1000 dilution
 Lysates/proteins at 20-50 µg per lane.
 Predicted band size: 48 kDa
 Observed band size: 54 kDa



Western blot-Anti-SMAD3 pAb

Lane 1: Mouse brain tissue lysate
 Lane 2: Mouse heart tissue lysate
 Lane 3: Rat kidney tissue lysate
 Lane 4: Rat liver tissue lysate
 All lanes: Anti-SMAD3 at 1:1000 dilution
 Lysates/proteins at 20-50 µg per lane.
 Predicted band size: 48 kDa
 Observed band size: 54 kDa



Immunohistochemistry-Anti-SMAD3 pAb


Immunohistochemical analysis of paraffin-embedded human pancreatic cancer using anti-SMAD3 Rabbit Antibody at 1:100 dilution. Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

Anti-SMAD3 Rabbit pAb

WL02288


For Research Use Only. Not For Use In Diagnostic Procedures

Product Information




Immunohistochemistry-Anti-SMAD3 pAb

Immunohistochemical analysis of paraffin-embedded human breast cancer using anti-SMAD3 Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunohistochemistry-Anti-SMAD3 pAb

Immunohistochemical analysis of paraffin-embedded rat kidney using anti-SMAD3 Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunofluorescence-Anti-SMAD3 pAb

Immunofluorescence analysis of paraffin-embedded mouse kidney using anti-SMAD3 Rabbit Antibody at 1:300 dilution.
Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0