

Anti-MCK10/NEP Rabbit pAb



WL05063

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-MCK10/NEP Rabbit pAb	
Source	Rabbit	
Species reactivity	Human, Mouse, Rat	
Tested applications	WB	1:500-1:1000
Pack size	50/100/200/500/1000µl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide	

General Information

Background Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. It belongs to a subfamily of tyrosine kinase receptors with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain, and that are activated by various types of collagen.

Immunogen Polyclonal antibody is produced by immunizing animals with a synthetic peptide of MCK10/NEP.

Purification Polyclonal antibody was purified by immunogen affinity chromatography.


Anti-MCK10/NEP Rabbit pAb



WL05063


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images



Western blot-Anti-MCK10/NEP pAb

Lane 1: Human HeLa cell lysate
 Lane 2: Human MCF-7 cell lysate
 Lane 3: Human HEK-293 cell lysate
 All lanes: Anti-MCK10/NEP at 1:1000 dilution
 Lysates/proteins at 20-50 µg per lane.
 Predicted band size: 101 kDa
 Observed band size: 101 kDa



Western blot-Anti-MCK10/NEP pAb

Lane 1: Mouse liver tissue lysate
 Lane 2: Rat stomach tissue lysate
 All lanes: Anti-MCK10/NEP at 1:1000 dilution
 Lysates/proteins at 20-50 µg per lane.
 Predicted band size: 101 kDa
 Observed band size: 101 kDa