Product Datasheet



Anti-Cyclin A2 Rabbit pAb

WL02964

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name Anti-Cyclin A2 Rabbit pAb

Source Rabbit

Species reactivity Human, Mouse, Rat

Tested applications WB 1:500-1:1000

Pack size 50/100/200/500/1000μl

Storage Store at -20°C. Avoid freeze/thaw cycles.

 $\textbf{Storage buffer} \hspace{1.5cm} \textbf{Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 } \mu\text{g/ml} \\$

BSA, 50% glycerol and less than 0.02% sodium azide

General Information

Background While overcoming the G1/S checkpoint to commence DNA replication

requires cyclin E, and traversing the G2/M checkpoint to initiate mitosis requires cyclin B to be present, cyclin A seems to be required for both S-

phase and M-phase. A number of studies have described the ability of overexpressed cyclin A to accellerate the G1 to S transition causing DNA

replication, and cyclin A antisense DNA can prevent DNA replication. Cyclin A availability is apparently the rate-limiting step for entry into mitosis, and cyclin A is required for completion of prophase. At late prophase, cyclin A

may no longer be necessary as cdc2/cyclinB1 becomes active.

Immunogen Polyclonal antibody is produced by immunizing animals with a synthetic

peptide of Cyclin A2.

Purification Polyclonal antibody was purified by immunogen affinity chromatography.

Product Datasheet

Wanleibio




Anti-Cyclin A2 Rabbit pAb

WL02964


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images



Western blot-Anti-Cyclin A2 pAb

Lane 1: Human HepG2 cell lysate
Lane 2: Human Hela cell lysate
Lane 3: Human BGC-823 cell lysate
Lane 4: Human MGC-803 cell lysate
All lanes: Anti-Cyclin A2 at 1:1000 dilution
Lysates/proteins at 20-50 µg per lane.
Predicted band size: 49 kDa



Western blot-Anti-Cyclin A2 pAb

Observed band size: 49 kDa

Lane 1: Mouse kidney tissue lysate
Lane 2: Mouse heart tissue lysate
Lane 3: Rat liver tissue lysate

All lanes: Anti-Cyclin A2 at 1:1000 dilution Lysates/proteins at 20-50 µg per lane.

Predicted band size: 49 kDa Observed band size: 49 kDa

Wanleibio Co.,Ltd. 400-602-0407 sales@wanleibio.com www.wanleibio.com Wanleibio Co.,Ltd. 400-602-0407 sales@wanleibio.com www.wanleibio.com