

## Anti-Factor X Rabbit pAb



WL02873

For Research Use Only. Not For Use In Diagnostic Procedures

## Product Information

<b>Product name</b>	Anti-Factor X Rabbit pAb	
<b>Source</b>	Rabbit	
<b>Species reactivity</b>	Human, Mouse, Rat	
<b>Tested applications</b>	WB	1:500-1:1000
	IHC	1:100-1:400
	IF	1:200
<b>Cellular localization</b>	Secreted	
<b>Pack size</b>	50/100/200/500/1000μl	
<b>Storage</b>	Store at -20°C. <b>Avoid freeze/thaw cycles.</b>	
<b>Storage buffer</b>	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 μg/ml	
	BSA, 50% glycerol and less than 0.02% sodium azide	

## General Information

**Background**

Factor X, also known by the eponym Stuart–Prower factor, is an enzyme of the coagulation cascade. It is a serine endopeptidase. Factor X is synthesized in the liver and requires vitamin K for its synthesis. Factor X is activated, by hydrolysis, into factor Xa by both factor IX and factor VII with its cofactor, tissue factor. The mature form of Factor X (Factor Xa) is generated by Factor IXa- or Factor VIIa-mediated cleavage at the tripeptide sequence, Arg-Lys-Arg, to yield a disulfide linked dimer. Together with the cofactor Factor Va and  $\text{Ca}^{2+}$  on the surface of platelets or endothelial cells, Factor Xa coordinates as part of the prothrombinase complex, which mediates proteolysis of Prothrombin into active Thrombin.

**Immunogen**

Polyclonal antibody is produced by immunizing animals with a synthetic peptide of Factor X.

**Purification**

Polyclonal antibody was purified by immunogen affinity chromatography.


## Anti-Factor X Rabbit pAb



WL02873


For Research Use Only. Not For Use In Diagnostic Procedures

## Product Images



Western blot-Anti-Factor X pAb

Lane 1: Human plasma lysate  
**All lanes:** Anti-Factor X at 1:1000 dilution  
 Lysates/proteins at 20-50 μg per lane.  
 Predicted band size: 54 kDa  
 Observed band size: 74 kDa



Western blot-Anti-Factor X pAb

Lane 1: Mouse liver tissue lysate  
 Lane 2: Rat liver tissue lysate  
**All lanes:** Anti-Factor X at 1:1000 dilution  
 Lysates/proteins at 20-50 μg per lane.  
 Predicted band size: 54 kDa  
 Observed band size: 74 kDa