

Anti-NOS2/INOS Rabbit pAb



WL0992a

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-NOS2/INOS Rabbit pAb	
Source	Rabbit	
Species reactivity	Human, Mouse, Rat	
Tested applications	WB	1:500-1:1000
	IHC	1:100-1:300
Cellular localization	Cytoplasm	
Pack size	50/100/200/500/1000μl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 μg/ml	
	BSA, 50% glycerol and less than 0.02% sodium azide	

General Information

Background	Nitric oxide (NO) has a broad range of biological activities and has been implicated in signaling pathways in phylogenetically diverse species. Nitric oxide synthases (NOSs), the enzymes responsible for synthesis of NO, contain an N-terminal oxygenase domain and a C-terminal reductase domain. NOS activity requires homodimerization as well as three cosubstrates (L-arginine, NADPH and O ₂) and five cofactors or prosthetic groups. Several distinct NOS isoforms have been described and been shown to represent the products of three distinct genes. These include two constitutive Ca ²⁺ /CaM-dependent forms of NOS, including NOS1 whose activity was first identified in neurons, and NOS3, first identified in endothelial cells. The inducible form of NOS, NOS2 (also designated iNOS), is Ca ²⁺ -independent and is expressed in a broad range of cell types.
Immunogen	Polyclonal antibody is produced by immunizing animals with a synthetic peptide of NOS2/INOS.
Purification	Polyclonal antibody was purified by immunogen affinity chromatography.


Anti-NOS2/INOS Rabbit pAb



WL0992a


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Western blot-Anti-NOS2/INOS pAb

Lane 1: Human Hela cell lysate
 All lanes: Anti-NOS2/INOS at 1:1000 dilution
 Lysates/proteins at 20-50 μg per lane.
 Predicted band size: 131 kDa
 Observed band size: 130 kDa



Western blot-Anti-NOS2/INOS pAb

Lane 1: Mouse colon tissue lysate
 Lane 2: Rat liver tissue lysate
 All lanes: Anti-NOS2/INOS at 1:1000 dilution
 Lysates/proteins at 20-50 μg per lane.
 Predicted band size: 131 kDa
 Observed band size: 130 kDa



Immunohistochemistry-Anti-NOS2/INOS pAb

Immunohistochemical analysis of paraffin-embedded mouse kidney using anti-NOS2/INOS Rabbit Antibody at 1:200 dilution.
 Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0


Anti-NOS2/INOS Rabbit pAb



For Research Use Only. Not For Use In Diagnostic Procedures

WL0992a

Product Information



Immunohistochemistry-Anti-NOS2/INOS pAb

Immunohistochemical analysis of paraffin-embedded mouse lung using anti-NOS2/INOS Rabbit Antibody at 1:300 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0