## **Product Datasheet**

# Wanleib

## Anti-NQO1 Rabbit pAb

WL04860

For Research Use Only. Not For Use In Diagnostic Procedures

## **Product Information**

**Product name** Anti-NQO1 Rabbit pAb

Source Rabbit

**Species reactivity** Human, Mouse, Rat

**Tested applications** WB 1:500-1:1000

> IHC 1:100-1:400

> ΙF 1:100-1:400

**Cellular localization** Cytoplasm

50/100/200/500/1000µl Pack size

Store at -20°C. Avoid freeze/thaw cycles. Storage

Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml Storage buffer

BSA, 50% glycerol and less than 0.02% sodium azide

## **General Information**

**Background** NQO1, also named as DTD , QR1, belongs to the NAD(P)H dehydrogenase

> (quinone) family. This enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinons involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in

prothrombin synthesis.

**Immunogen** Polyclonal antibody is produced by immunizing animals with a synthetic

peptide of NQO1.

Purification Polyclonal antibody was purified by immunogen affinity chromatography.

## **Product Datasheet**


## Anti-NQO1 Rabbit pAb



WL04860


For Research Use Only. Not For Use In Diagnostic Procedures

## **Product Images**




#### Western blot-Anti-NQO1 pAb

Lane 1: Human HUVEC cell lysate Lane 2: Human SW480 cell lysate Lane 3: Human A549 cell lysate All lanes: Anti-NQO1 at 1:1000 dilution Lysates/proteins at 20-50 µg per lane. Predicted band size: 31 kDa Observed band size: 30 kDa



#### Western blot-Anti-NQO1 pAb

Lane 1: Mouse liver tissue lysate Lane 2: Rat colon tissue lysate All lanes: Anti-NQO1 at 1:1000 dilution Lysates/proteins at 20-50 µg per lane. Predicted band size: 31 kDa Observed band size: 30 kDa



#### Immunohistochemistry-Anti-NQO1 pAb

Immunohistochemical analysis of paraffin-embedded mouse kidney using anti-NQO1 Rabbit Antibody at 1:200 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

Wanleibio Co., Ltd. 400-602-0407 sales@wanleibio.com www.wanleibio.com Wanleibio Co., Ltd. 400-602-0407 www.wanleibio.com sales@wanleibio.com


## Anti-NQO1 Rabbit pAb



WL04860

For Research Use Only. Not For Use In Diagnostic Procedures

## **Product Information**



## Immunohistochemistry-Anti-NQO1 pAb

Immunohistochemical analysis of paraffin-embedded mouse kidney using anti-NQO1 Rabbit Antibody at 1:200 dilution.


Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



### Immunohistochemistry-Anti-NQO1 pAb

Immun ohist ochemical analysis of paraffin-embedded mouse liver using anti-NQO1 Rabbit Antibody at 1:200 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



#### Immunohistochemistry-Anti-NQO1 pAb

Immun ohist ochemical analysis of paraffin-embedded mouse liver using anti-NQO1 Rabbit Antibody at 1:200 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

## **Product Datasheet**


## Wanleibio 🖦

## Anti-NQO1 Rabbit pAb

WL04860

For Research Use Only. Not For Use In Diagnostic Procedures

## **Product Images**



#### Immunofluorescence-Anti-NQO1 pAb

 $Immun of luorescence\ analysis\ of\ paraffin-embedded\ rat\ kidney\ using\ anti-NQO1\ Rabbit\ Antibody\ at\ 1:200\ dilution.$ 

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

Wanleibio Co.,Ltd. 400-602-0407 sales@wanleibio.com www.wanleibio.com Wanleibio Co.,Ltd. 400-602-0407 sales@wanleibio.com www.wanleibio.com