# Wanleibio

# Anti-FGF-1 Rabbit pAb

WLH4209

For Research Use Only.Not For Use In Diagnostic Procedures

### **Product Information**

Product name	Anti-FGF-1 Rabbit pAb		
Source	Rabbit		
Species reactivity	Human, Mouse, Rat		
Tested applications	IHC, IF	1:100-1:500	
Cellularlocalization	Secreted. Cytoplasm		
Packsize	50/100/200/500/1000µl		
Storage	Store at -20°C. Avoid freeze/thaw cycles.		
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 $\mu g/ml$		
	BSA, 50% glycerol and less than 0.02% sodium azide		

## Product Datasheet

# Anti-FGF-1 Rabbit pAb



WLH4209

For Research Use Only.Not For Use In Diagnostic Procedures

#### Product Images



#### Immunohistochemistry-Anti-FGF-1 pAb

Immunohistochemical analysis of paraffin-embedded mouse brain using anti-FGF-1 Rabbit Antibody at 1:150 dilution. Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



#### Immunohistochemistry-Anti-FGF-1 pAb

Immunohistochemical analysis of paraffin-embedded rat kidney using anti-FGF-1 Rabbit Antibody at 1:150 dilution. Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

# **General Information**

and invasion.	development, cell growth, morphogene	members possess broad mitogenic and cell survival activities, and are		c FGF, are feration of cells GF family ties, and are bryonic
involved in a variety of biological processes, including embryonic	members possess broad mitogenic and o			GF family
involved in a variety of biological processes, including embryonic		of mesenchymal, epithelial and neuroectodermal origin. FGF family	BuckBrowning Contraction Contraction	feration of ce
of mesenchymal, epithelial and neuroectodermal origin. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic	of mesenchymal, epithelial and neuroed	, , , , , , , , , , , , , , , , , , , ,	Background Fibroblast growth factor-1 (FGF-1), also designated acidic F	c FGF, are
members possess broad mitogenic and cell survival activities, and are	members of a family of growth factors th of mesenchymal, epithelial and neuroed	members of a family of growth factors that stimulate proliferation of ce		FGF, and

ImmunogenPolyclonal antibody is produced by immunizing animals with a synthetic<br/>peptide of FGF-1.PurificationPolyclonal antibody was purified by immunogen affinity chromatography.

# Immu using a Perfor

#### Immunofluorescence-Anti-FGF-1 pAb

Immunofluorescence analysis of paraffin-embedded human kidney cancer using anti-FGF-1 Rabbit Antibody at 1:200 dilution. Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0