Wanleibio

WL03211

Anti-LRP1 Rabbit pAb

For Research Use Only.Not For Use In Diagnostic Procedures

Product Information

General Information

Background

Immunogen

Product name	Anti-LRP1 Rabbit pAb	
Source	Rabbit	
Species reactivity	Human, Mouse, Rat	
Tested applications	WB	1:1000-1:2000
	IHC	1:150
	IF	1:500
Cellularlocalization	Cytoplasm. Nucleus	
Pack size	50/100/200/500/1000µl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 $\mu g/ml$	
	BSA, 50% glycerol and less than 0.02% sodium azide	


Product Datasheet

Anti-LRP1 Rabbit pAb

Wanleibio


For Research Use Only.Not For Use In Diagnostic Procedures

Product Images




Western blot-Anti-LRP1 pAb

Lane 1: Human HepG2 cell lysate Lane 2: Human Hela cell lysate Lane 3: Human SH-SY5Y cell lysate All lanes: Anti-LRP1 at 1:1000 dilution Lysates/proteins at 20-50 µg per lane. Predicted band size: 505 kDa Observed band size: 85-90 kDa



Western blot-Anti-LRP1 pAb

Lane 1: Mouse lung tissue lysate Lane 2: Mouse liver tissue lysate Lane 3: Rat liver tissue lysate Lane 4: Rat lung tissue lysate All lanes: Anti-LRP1 at 1:1000 dilution Lysates/proteins at 20-50 µg per lane. Predicted band size: 505 kDa Observed band size: 85-90 kDa



Immunohistochemistry-Anti-LRP1 pAb

Immunohistochemical analysis of paraffin-embedded mouse lung using anti-LRP1 Rabbit Antibody at 1:150 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

Purification	

Polyclonal antibody was purified by Protein A affinity chromatography.

LRP1 encodes a member of the low-density lipoprotein receptor family of

proteins. The encoded preproprotein is proteolytically processed by furin

implicated in Alzheimer's disease (AD), as the human LRP gene localizes

Polyclonal antibody is produced by immunizing animals with a synthetic

to generate 515 kDa and 85 kDa subunits that form the mature receptor. Proper folding and trafficking of LRP1 is facilitated by the receptorassociated protein (RAP), a molecular chaperone. The uptake of all known

ligands through LRP1 can be blocked by RAP, which induces a conformational change in the receptor that renders it unable to bind

ligands. LRP1, which is expressed in brain, liver and lung, is also

to a potential AD locus on chromosome 12.

peptide of LRP1.

WL03211


Wanleibio

WL03211

Anti-LRP1 Rabbit pAb

For Research Use Only.Not For Use In Diagnostic Procedures

Product Information



Immunofluorescence-Anti-LRP1 pAb

Immunofluorescence analysis of paraffin-embedded mouse lung using anti-LRP1 Rabbit Antibody at 1:500 dilution. Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

Wanleibio Co.,Ltd. 400-602-0407 sales@wanleibio.com www.wanleibio.com