# Wanleibio

WL02989

# Anti-p-TrkA (Y757) Rabbit pAb

For Research Use Only.Not For Use In Diagnostic Procedures

### **Product Information**

Product name	Anti-p-TrkA (Y757) Rabbit pAb	
Source	Rabbit	
Species reactivity	Human, Mouse, Rat	
Tested applications	Western blot	1:1000-1:2000
	*Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own experiment using appropriate negative and positive controls.	
Pack size	50/100/200/500/1000µl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 $\mu g/ml$	
	BSA, 50% glycerol and less than 0.02% sodium azide	

## Product Datasheet

# Anti-p-TrkA (Y757) Rabbit pAb

For Research Use Only. Not For Use In Diagnostic Procedures





#### Western blot-Anti-p-TrkA (Y757) pAb

Lane 1: Human HepG2 cell lysate Lane 2: Human Hela cell lysate Lane 3: Human BGC-823 cell lysate Lane 4: Human MGC-803 cell lysate All lanes: Anti-p-TrkA (Y757) at 1:1000 dilution Lysates/proteins at 20-50 µg per lane. Predicted band size: 87 kDa Observed band size: 87/140 kDa



#### Western blot-Anti-p-TrkA (Y757) pAb

Lane 1: Mouse kidney tissue lysate Lane 2: Mouse heart tissue lysate Lane 3: Rat liver tissue lysate Lane 4: Rat brain tissue lysate All lanes: Anti-p-TrkA (Y757) at 1:1000 dilution Lysates/proteins at 20-50 µg per lane. Predicted band size: 87 kDa Observed band size: 87/140 kDa

## General Information

BackgroundTrkA is the high affinity catalytic receptor for the neurotrophin, Nerve<br/>Growth Factor, or "NGF". As such, it mediates the multiple effects of NGF,<br/>which include neuronal differentiation and avoidance of programmed cell<br/>death. TrkA was originally cloned from a colon tumor; the cancer occurred<br/>via a translocation, which resulted in the activation of the TrkA kinase<br/>domain. However, TrkA itself does not appear to be an oncogene.<br/>Phosphorylation at Y757 site reflects TrkA kinase activity.

Immunogen	Polyclonal antibody is produced by immunizing animals with a synthetic peptide of p-TrkA (Y757).
Purification	Polyclonal antibody was purified by Protein A affinity chromatography.



WL02989