### **Product Datasheet**

# Wanleibio 🖦

## Anti-Apolipoprotein CIII Rabbit pAb

WL02058

For Research Use Only. Not For Use In Diagnostic Procedures

### **Product Information**

Product name Anti-Apolipoprotein CIII Rabbit pAb

Source Rabbit

Species reactivity Human, Mouse, Rat

Tested applications WB 1:1000-1:2000

IHC 1:100

IF 1:100-1:200

**Cellular localization** Secreted

Pack size 50/100/200/500/1000μl

Storage Store at -20°C. Avoid freeze/thaw cycles.

Storage buffer Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml

BSA, 50% glycerol and less than 0.02% sodium azide

### **General Information**

Background Apolipoprotein C-III is a very low density lipoprotein (VLDL) protein.

Apolipoproteins are protein components of plasma lipoproteins. The apolipoprotein C gene family encodes four homologous proteins designated apoC-I to -IV, which specifically modulate the metabolism of triglyceride-rich lipoproteins. The human apoC-III gene maps to chromosome 11q23 and encodes a protein that may delay catabolism of triglyceride-rich particles by inhibiting lipoprotein lipase and hepatic lipase. APOC3 has 99 amino acids and T64,T94,S95 are glycosylated.An

increase in apoC-III levels induces the development of

hypertriglyceridemia.

Immunogen Polyclonal antibody is produced by immunizing animals with a synthetic

peptide of Apolipoprotein CIII.

Purification Polyclonal antibody was purified by Protein A affinity chromatography.

### **Product Datasheet**

**Product Images** 

# Anti-Apolipoprotein CIII Rabbit pAb





WL02058

### For Research Use Only. Not For Use In Diagnostic Procedures

# kDa 1 17-10-


### Western blot-Anti-Apolipoprotein CIII pAb

Lane 1: Human HEP-G2 cell lysate

All lanes: Anti-Apolipoprotein CIII at 1:1000 dilution

Lysates/proteins at 20-50 µg per lane.

Predicted band size: 11 kDa Observed band size: 11 kDa



#### Western blot-Anti-Apolipoprotein CIII pAb


Lane 1: Mouse liver tissue lysate

Lane 2: Rat liver tissue lysate

All lanes: Anti-Apolipoprotein CIII at 1:1000 dilution

Lysates/proteins at 20-50 µg per lane.

Predicted band size: 11 kDa Observed band size: 11 kDa



#### Immunohistochemistry-Anti-Apolipoprotein CIII pAb

Immunohistochemical analysis of paraffin-embedded rat abnormal liver using anti-Apolipoprotein CIII Rabbit Antibody at 1:100 dilution. Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

Wanleibio Co., Ltd. 400-602-0407 sales@wanleibio.com www.wanleibio.com Wanleibio Co., Ltd. 400-602-0407 sales@wanleibio.com www.wanleibio.com


## Anti-Apolipoprotein CIII Rabbit pAb



WL02058


For Research Use Only. Not For Use In Diagnostic Procedures

### **Product Information**



### Immunohistochemistry-Anti-Apolipoprotein CIII pAb

Immunohistochemical analysis of paraffin-embedded human pancreatic cancer using anti-Apolipoprotein CIII Rabbit Antibody at 1:100 dilution. Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



### Immunofluorescence-Anti-Apolipoprotein CIII pAb

Immunofluorescence analysis of paraffin-embedded mouse intestine using anti-Apolipoprotein CIII Rabbit Antibody at 1:100 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

www.wanleibio.com