

Anti-CDK6 Rabbit pAb



WL03460

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

| | | |
|------------------------------|--|--------------|
| Product name | Anti-CDK6 Rabbit pAb | |
| Source | Rabbit | |
| Species reactivity | Human, Mouse, Rat | |
| Tested applications | WB | 1:500-1:1000 |
| | IHC | 1:200 |
| | IF | 1:200 |
| Cellular localization | Cell projection. Cytoplasm. Cytoskeleton. Nucleus. | |
| Pack size | 50/100/200/500/1000μl | |
| Storage | Store at -20°C. Avoid freeze/thaw cycles. | |
| Storage buffer | Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 μg/ml | |
| | BSA, 50% glycerol and less than 0.02% sodium azide | |

General Information

| | |
|---------------------|---|
| Background | The cyclin-dependent kinases form complexes with their cyclin partners and with CDK inhibitors. CDK6 and CDK4 associate with the D-type cyclins and target the retinoblastoma protein, allowing passage through the G1/S phase restriction point. In cervical cancer cells, CDK6 function has been shown to be altered indirectly by the p16 inhibitor. CDK6 is also overexpressed in tumors that exhibit drug resistance, for example glioma malignancies exhibit resistance to chemotherapy using temozolomide (TMZ) when they have a mutation overexpressing CDK6. |
| Immunogen | Polyclonal antibody is produced by immunizing animals with a synthetic peptide of CDK6. |
| Purification | Polyclonal antibody was purified by immunogen affinity chromatography. |


Anti-CDK6 Rabbit pAb



WL03460


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Western blot-Anti-CDK6 pAb

Lane 1: Human Hela cell lysate
 Lane 2: Human HL-60 cell lysate
 All lanes: Anti-CDK6 at 1:1000 dilution
 Lysates/proteins at 20-50 μg per lane.
 Predicted band size: 37 kDa
 Observed band size: 37 kDa



Western blot-Anti-CDK6 pAb

Lane 1: Mouse lung tissue lysate
 Lane 2: Rat kidney tissue lysate
 All lanes: Anti-CDK6 at 1:1000 dilution
 Lysates/proteins at 20-50 μg per lane.
 Predicted band size: 37 kDa
 Observed band size: 37 kDa



Immunohistochemistry-Anti-CDK6 pAb

Immunohistochemical analysis of paraffin-embedded human rectum cancer using anti-CDK6 Rabbit Antibody at 1:200 dilution. Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0


Anti-CDK6 Rabbit pAb



WL03460

For Research Use Only. Not For Use In Diagnostic Procedures


Product Information



Immunohistochemistry-Anti-CDK6 pAb

Immunohistochemical analysis of paraffin-embedded mouse brain using anti-CDK6 Rabbit Antibody at 1:200 dilution.


Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunofluorescence-Anti-CDK6 pAb

Immunofluorescence analysis of paraffin-embedded human colon using anti-CDK6 Rabbit Antibody at 1:200 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0



Immunofluorescence-Anti-CDK6 pAb

Immunofluorescence analysis of paraffin-embedded mouse pancreas using anti-CDK6 Rabbit Antibody at 1:200 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0