

Anti-p62 Rabbit pAb



WL02385

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-p62 Rabbit pAb		
Source	Rabbit		
Species reactivity	Human, Mouse, Rat, Pig		
Tested applications	Western blot	1:500-1:1000	
	Immunohistochemistry	1:100-1:300	
<i>*Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own experiment using appropriate negative and positive controls.</i>			
Molecular Wt.	62 kDa		
Pack size	50/100/200/500/1000μl		
Storage	Store at -20°C. Avoid freeze/thaw cycles.		
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 μg/ml		
	BSA, 50% glycerol and less than 0.02% sodium azide		

General Information

Background	P62 gene encodes a multifunctional protein that binds ubiquitin and regulates activation of the nuclear factor kappa-B (NF-κB) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF-κB in response to upstream signals. Mutations in the ubiquitin-associated (UBA) domain of the Sequestosome 1 protein (SQSTM1), also designated p62 or ZIP, commonly cause Paget's disease since the UBA is necessary for aggregate sequestration and cell survival.
Immunogen	Polyclonal antibody is produced by immunizing animals with a synthetic peptide of p62.
Purification	Polyclonal antibody was purified by immunogen affinity chromatography.


Anti-p62 Rabbit pAb



WL02385


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Western blot-Anti-p62 pAb

Lane 1: Human HepG2 cell lysate 30μg
 Lane 2: Human Hela cell lysate 30μg
 Lane 3: Human BGC-823 cell lysate 30μg
Separation gel: 10% polyacrylamide
Electrophoresis: 100V, 4°C, 3h
Transmembrane: 100V, 4°C, 1h
Blocking: 5% w/v nonfat dry milk, 1× TBST, at RT with gentle shaking
Primary antibody: 1:1000 in blocking buffer, 4°C, overnight
Secondary antibody (WLA023a) : 1:5000-1:10000, 45min
Detection: ECL, 30s-2min



Western blot-Anti-p62 pAb

Lane 1: Mouse liver tissue lysate 30μg
 Lane 2: Rat lung tissue lysate 30μg
Separation gel: 10% polyacrylamide
Electrophoresis: 100V, 4°C, 3h
Transmembrane: 100V, 4°C, 1h
Blocking: 5% w/v nonfat dry milk, 1× TBST, at RT with gentle shaking
Primary antibody: 1:1000 in blocking buffer, 4°C, overnight
Secondary antibody (WLA023a) : 1:5000-1:10000, 45min
Detection: ECL, 30s-2min



Immunohistochemistry-Anti-p62 pAb

Sample: Human colon cancer tissue
Antigen retrieval: pH 9.0 Tris-EDTA buffer
Primary antibody: 1:200, 4°C, overnight
Secondary antibody-Biotin: 1:150, 37°C, 1h
Streptavidin-HRP: 1:200, 37°C, 30min
Color Developing: DAB


Anti-p62 Rabbit pAb



WL02385


For Research Use Only. Not For Use In Diagnostic Procedures

Product Information




Immunohistochemistry-Anti-p62 pAb

Sample: Human pancreatic cancer tissue
Antigen retrieval: pH 9.0 Tris-EDTA buffer
Primary antibody: 1:200, 4°C, overnight
Secondary antibody-Biotin: 1:150, 37°C, 1h
Streptavidin-HRP: 1:200, 37°C, 30min
Color Developing: DAB



Immunohistochemistry-Anti-p62 pAb

Sample: Mouse liver tissue
Antigen retrieval: pH 9.0 Tris-EDTA buffer
Primary antibody: 1:200, 4°C, overnight
Secondary antibody-Biotin: 1:150, 37°C, 1h
Streptavidin-HRP: 1:200, 37°C, 30min
Color Developing: DAB



Immunohistochemistry-Anti-p62 pAb

Sample: Mouse kidney tissue
Antigen retrieval: pH 9.0 Tris-EDTA buffer
Primary antibody: 1:200, 4°C, overnight
Secondary antibody-Biotin: 1:150, 37°C, 1h
Streptavidin-HRP: 1:200, 37°C, 30min
Color Developing: DAB


Anti-p62 Rabbit pAb



WL02385


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images




Immunohistochemistry-Anti-p62 pAb

Sample: Rat bladder tissue
Antigen retrieval: pH 9.0 Tris-EDTA buffer
Primary antibody: 1:200, 4°C, overnight
Secondary antibody-Biotin: 1:150, 37°C, 1h
Streptavidin-HRP: 1:200, 37°C, 30min
Color Developing: DAB



Immunohistochemistry-Anti-p62 pAb

Sample: Rat liver tissue
Antigen retrieval: pH 9.0 Tris-EDTA buffer
Primary antibody: 1:200, 4°C, overnight
Secondary antibody-Biotin: 1:150, 37°C, 1h
Streptavidin-HRP: 1:200, 37°C, 30min
Color Developing: DAB



Immunohistochemistry-Anti-p62 pAb

Sample: Rat kidney tissue
Antigen retrieval: pH 9.0 Tris-EDTA buffer
Primary antibody: 1:200, 4°C, overnight
Secondary antibody-Biotin: 1:150, 37°C, 1h
Streptavidin-HRP: 1:200, 37°C, 30min
Color Developing: DAB


Anti-p62 Rabbit pAb



WL02385

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information



Immunohistochemistry-Anti-p62 pAb

Sample: Rat uterus tissue

Antigen retrieval: pH 9.0 Tris-EDTA buffer

Primary antibody: 1:200, 4°C, overnight

Secondary antibody-Biotin: 1:150, 37°C, 1h

Streptavidin-HRP: 1:200, 37°C, 30min

Color Developing: DAB