

Anti-Podoplanin Rabbit pAb



WLH4439

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-Podoplanin Rabbit pAb	
Source	Rabbit	
Species reactivity	Human, Mouse, Rat	
Tested applications	WB	1:500-1:1000
	IHC	1:100-1:400
Cellular localization	Cell junction, Cell membrane, Cell projection, Cytoplasm, Membrane.	
Pack size	50/100/200/500/1000µl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide	

General Information

Background Podoplanin is type-I integral membrane glycoprotein with diverse distribution in human tissues. The physiological function of this protein may be related to its mucin-type character. The homologous protein in other species has been described as a differentiation antigen and influenza-virus receptor. The specific function of this protein has not been determined but it has been proposed as a marker of lung injury.

Immunogen Polyclonal antibody is produced by immunizing animals with a synthetic peptide of Podoplanin.

Purification Polyclonal antibody was purified by immunogen affinity chromatography.


Anti-Podoplanin Rabbit pAb



WLH4439

For Research Use Only. Not For Use In Diagnostic Procedures

Product Images



Western blot-Anti-Podoplanin pAb

Lane 1: Human HCT116 cell lysate

Lane 2: Human TPC-1 cell lysate


Lane 3: Rat RSC96 cell lysate

All lanes: Anti-Podoplanin at 1:1000 dilution

Lysates/proteins at 20-50 µg per lane.

Predicted band size: 17 kDa

Observed band size: 36-43 kDa



Immunohistochemistry-Anti-Bag3 pAb

Sample: Mouse kidney tissue


Antigen retrieval: pH 9.0 Tris-EDTA buffer

Primary antibody: 1:200, 4°C, overnight

Secondary antibody-Biotin: 1:150, 37°C, 1h

Streptavidin-HRP: 1:200, 37°C, 30min

Color Developing: DAB



Immunohistochemistry-Anti-Bag3 pAb

Sample: Rat kidney tissue

Antigen retrieval: pH 9.0 Tris-EDTA buffer

Primary antibody: 1:200, 4°C, overnight

Secondary antibody-Biotin: 1:150, 37°C, 1h

Streptavidin-HRP: 1:200, 37°C, 30min

Color Developing: DAB


Anti-Podoplanin Rabbit pAb



WLH4439

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information



Immunohistochemistry-Anti-Bag3 pAb

Sample: Rat kidney tissue

Antigen retrieval: pH 9.0 Tris-EDTA buffer

Primary antibody: 1:200, 4°C, overnight

Secondary antibody-Biotin: 1:150, 37°C, 1h

Streptavidin-HRP: 1:200, 37°C, 30min

Color Developing: DAB