

Anti-p62 Rabbit pAb



WLH4306

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-p62 Rabbit pAb	
Source	Rabbit	
Species reactivity	Human, Mouse, Rat, Rabbit	
Tested applications	IHC, IF	1:200-1:400
Cellular localization	Cytoplasm. Late endosome. Nucleus.	
Pack size	50/100/200/500/1000µl	
Storage	Store at -20°C. Avoid freeze/thaw cycles.	
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide	

General Information

Background	P62 gene encodes a multifunctional protein that binds ubiquitin and regulates activation of the nuclear factor kappa-B (NF-κB) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF-κB in response to upstream signals. Mutations in the ubiquitin-associated (UBA) domain of the Sequestosome 1 protein (SQSTM1), also designated p62 or ZIP, commonly cause Paget's disease since the UBA is necessary for aggregate sequestration and cell survival.
Immunogen	Polyclonal antibody is produced by immunizing animals with a recombinant protein of p62.
Purification	Polyclonal antibody was purified by immunogen affinity chromatography.


Anti-p62 Rabbit pAb




WLH4306

For Research Use Only. Not For Use In Diagnostic Procedures


Product Images

**Immunohistochemistry-Anti-p62 pAb**

Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue using anti-p62 Rabbit Antibody at 1:200 dilution. Perform heat mediated antigen retrieval with citrate buffer pH 6.0

**Immunohistochemistry-Anti-p62 pAb**

Immunohistochemical analysis of paraffin-embedded mouse spleen tissue using anti-p62 Rabbit Antibody at 1:200 dilution. Perform heat mediated antigen retrieval with citrate buffer pH 6.0

**Immunohistochemistry-Anti-p62 pAb**

Immunohistochemical analysis of paraffin-embedded rat liver tissue using anti-p62 Rabbit Antibody at 1:200 dilution. Perform heat mediated antigen retrieval with citrate buffer pH 6.0


Anti-p62 Rabbit pAb



WLH4306


For Research Use Only. Not For Use In Diagnostic Procedures

Product Information




Immunohistochemistry-Anti-p62 pAb

Immunohistochemical analysis of paraffin-embedded rat skin tissue using anti-p62 Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6.0



Immunohistochemistry-Anti-p62 pAb

Immunohistochemical analysis of paraffin-embedded rat stomach tissue using anti-p62 Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6.0



Immunohistochemistry-Anti-P62 pAb

Immunohistochemical analysis of paraffin-embedded rabbit spleen tissue using anti-p62 Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6.0


Anti-p62 Rabbit pAb



WLH4306

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information



Immunofluorescence-Anti-p62 pAb

Immunofluorescence analysis of paraffin-embedded human liver cancer using anti-p62 Rabbit Antibody at 1:200 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6.0