

Anti-BCL-10 Rabbit pAb



WLH4556

For Research Use Only. Not For Use In Diagnostic Procedures

Product Information

Product name	Anti-BCL-10 Rabbit pAb
Source	Rabbit
Species reactivity	Mouse, Rat
Tested applications	IHC 1:100-1:400
Cellular localization	Secreted and Cell membrane
Pack size	50/100/200/500/1000µl
Storage	Store at -20°C. Avoid freeze/thaw cycles.
Storage buffer	Supplied in 20 mM phosphate (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide

General Information

Background The was identified by its translocation in a case of mucosa-associated lymphoid tissue (MALT) lymphoma. BCL10 contains a caspase recruitment domain (CARD), and has been shown to induce apoptosis and to activate NF-kappaB.

Immunogen Polyclonal antibody is produced by immunizing animals with a synthetic peptide of BCL-10.

Purification Polyclonal antibody was purified by immunogen affinity chromatography.


Anti-BCL-10 Rabbit pAb



WLH4556


For Research Use Only. Not For Use In Diagnostic Procedures

Product Images

**Immunohistochemistry-Anti-BCL-10 pAb**


Immunohistochemical analysis of paraffin-embedded mouse ovary using anti-BCL-10 Rabbit Antibody at 1:200 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

**Immunohistochemistry-Anti-BCL-10 pAb**

Immunohistochemical analysis of paraffin-embedded rat ovary using anti-BCL-10 Rabbit Antibody at 1:200 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0

**Immunohistochemistry-Anti-BCL-10 pAb**

Immunohistochemical analysis of paraffin-embedded rat kidney using anti-BCL-10 Rabbit Antibody at 1:200 dilution.

Perform heat mediated antigen retrieval with Tris-EDTA buffer pH 9.0